

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 1996		3. REPORT TYPE AND DATES COVERED Professional Paper	
4. TITLE AND SUBTITLE ALGEBRAIC REPRESENTATIONS OF LINGUISTIC AND NUMERICAL MODIFICATIONS OF PROBABILITY STATEMENTS AND INFERENCES BASED ON A PRODUCT SPACE CONSTRUCTION				5. FUNDING NUMBERS PR: CD32 PE: 0601152N WU: DN300173	
6. AUTHOR(S) I. R. Goodman					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Command, Control and Ocean Surveillance Center (NCCOSC) RDT&E Division San Diego, CA 92152-5001				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Independent Research Program (IR) 800 North Quincy Street Arlington, VA 22217-5660				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.				12b. DISTRIBUTION CODE  19960404 087	
13. ABSTRACT (Maximum 200 words)  This paper addresses the problem of obtaining for any given probability space $(\Omega, B, P)$ an extension to a probability space $(\Omega_0, B_0, P_0)$ such that for any given function $f: [0,1]^n \rightarrow [0,1]$ , suitably analytic, and any given finite collection of ordinary events $a, b, c, \dots, m$ in sample space $B$ , with possible relations among them, there is an event $\alpha_f = \alpha_f(a, b, c, \dots, m)$ in $B_0$ (not dependent upon $P$ ) such that for all possible choices of $P$ and $a, b, c, \dots, m$ , $P_0(\alpha_f) = f(P(a), P(b), P(c), \dots, P(m)),$ provided the right-hand side is well-defined relative to $f$ . It is seen that the above procedure generalizes that of the problem of constructing "conditional event algebras" relative to all conditional probabilities. Applications to problems on data fusion and combination of evidence are also provided.  Published in <i>Proceedings of 32nd Annual Bayesian Research Conference</i> , February 17, 1994.					
14. SUBJECT TERMS Mission Area: Command and Control (C <sup>2</sup> ), Intelligence Probability Space Data Fusion				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT SAME AS REPORT		

UNCLASSIFIED

21a. NAME OF RESPONSIBLE INDIVIDUAL I. R. Goodman	21b. TELEPHONE <i>(include Area Code)</i> (619) 553-4014	21c. OFFICE SYMBOL Code 4221

## 32<sup>ND</sup> ANNUAL BAYESIAN RESEARCH CONFERENCE

# ALGEBRAIC REPRESENTATIONS OF LINGUISTIC AND NUMERICAL MODIFICATIONS OF PROBABILITY STATEMENTS AND INFERENCES BASED ON A PRODUCT SPACE CONSTRUCTION

Dr. I.R. Goodman,

NCCOSC RDTE DIV 4221  
53140 SYSTEMS ST ROOM 230  
SAN DIEGO, CA 92152-7550

Tel; (619) 553-4014

Fax: (619) 553-5793

Email: goodman@nosc.mil

### EXTENDED ABSTRACT

#### S u m m a r y

In brief, this paper addresses the problem of obtaining for any given probability space  $(\Omega, B, P)$  an extension to a probability space  $(\Omega_0, B_0, P_0)$  such that for any given function  $f: [0,1]^n \rightarrow [0,1]$ , suitably analytic, and any given finite collection of ordinary events  $a, b, c, \dots, m$  in sample space  $B$ , with possible relations among them, there is an event  $\alpha_f = \alpha_f(a, b, c, \dots, m)$  in  $B_0$  (not dependent upon  $P$ ) such that for all possible choices of  $P$  and  $a, b, c, \dots, m$ ,

$$P_0(\alpha_f) = f(P(a), P(b), P(c), \dots, P(m)) , \quad (1)$$

provided the right-hand side is well-defined relative to  $f$ . It is seen that the above procedure generalizes that of the problem of constructing "conditional event algebras" relative to all conditional probabilities. Applications to problems of data fusion and combination of evidence are also provided.

#### Motivations and More Details

Numerical modifications of individual probability expressions arise quite often based solely on probability evaluations themselves, as. e.g., (1), in the expansion  $f(P(a \bullet b), P(b)) = P(a \vee b') = 1 - P(b) + P(a \bullet b)$  or, (2), when  $a$  and  $b$  are  $P$ -independent,  $f(P(a), P(b)) = P(a \bullet b) = P(a) \cdot P(b)$ , where we note the first expression is an affine combination of  $P(b)$  and  $P(a \bullet b)$ , while the second expression is an arithmetic product of  $P(a)$  and  $P(b)$ . They can also arise when: (3), we wish to obtain weighted averages of calculated probabilities as in ,  $f(P(a), P(b)) = w \cdot P(a) + (1-w) \cdot P(b)$ , or perhaps (4), as an arithmetic division of  $P(a \bullet b)$  by  $P(b)$ , as in  $f(P(a \bullet b), P(b)) = P(a \bullet b) / P(b)$ , which of course we recognize as  $P(a|b)$ , the

conditional probability of a given b. Further modifications to previously calculated probabilities can occur when: (5), linguistic changes are imposed as in the modeling of (meta) statements such as  $s = P(b)$ , where  $b = \text{"It is very probable that } a\text{"}$ , where  $a = \text{"John will arrive tomorrow"}$  and where by prior knowledge we can estimate  $P(a)$ . Also, by prior knowledge, we conclude that the hedge "very probable" typically produces an exponentiation, of say 1.5, so that we obtain the estimate  $f(P(a)) = P(a)^{1.5}$ , quite reminiscent of the well-known use of exponential or other hedges in fuzzy set modeling. Similarly, (6), we may wish to obtain more complicated compounds of known probabilities in modeling empirically outputs of physical systems as, e.g., in the relation  $f(P(a), P(a \bullet b), P(b)) = P(a)^{2.6} + ((3.2) \cdot P(b) \cdot P(a \bullet b)) / (4 \cdot P(b) + P(a))$ .

Except for the first two examples, in general for each of the remaining examples 3-6, there is *no* corresponding event  $\alpha_f$ , some function (boolean or otherwise) of  $a, b, c, \dots$  in the original sample space  $B$  of the basic events  $a, b, c, \dots$  such that equation (1) is satisfied.

Thus, in general we cannot obtain probability evaluations of further logical combinations of expressions whose modified probability evaluations are known as in (1), such as in the obtaining of

$$P(\alpha \vee \beta) = P(\alpha) + P(\beta) - P(\alpha \bullet \beta) = s + t - P(\alpha \bullet \beta), \quad (2)$$

where analogous to (1)

$$P(\alpha) = s, \quad P(\beta) = t, \quad (3)$$

$s, t$  known functions of  $P(a), P(b), P(a \bullet b), P(c)$ , etc.

On the other hand, a remedy has been developed for the type of problem arising in the fourth example: arithmetic division of probabilities resulting in conditional probabilities can now be treated via some choice of a conditional event algebra (out of many possible candidates). Even though in this case  $\alpha$  and  $\beta$  in eq.(3) (where here  $s$  and  $t$  are arithmetic divisions representing conditional probabilities) do not lie in  $B$ , they do lie in a space naturally extending  $B$  - the conditional event algebra of choice. (See [1] for background and more details.)

In fact, a recent result [2] has indicated that the *only* mathematically desirable choice for a conditional event algebra is the one arising uniquely from the construction of the product space that has a countable infinity of independent marginal probability spaces, each identical to the one representing the original sample space of ordinary unconditional events [3]. In that space, typically, conditional events as  $\alpha_f$  satisfying eq.(1) for Example 4, where

$$f(P(a \bullet b), P(b)) = P(a \bullet b) / P(b) = P(alb), \quad (4)$$

take the form of the natural disjoint cartesian product counterpart to the expansion of arithmetic division in terms of an infinite power series:

$$\alpha_f = (alb) = \sum_{j=0}^{+\infty} (b' \times \dots \times b') \times (a \bullet b) \times \Omega \times \Omega \times \dots, \quad (5)$$

(j factors)

where  $\Omega$  is the universal event in sample space (sigma-algebra)  $B$ . Thus, for any given probability measure  $P$  over  $B$ , letting  $P_0$  denote the corresponding product space probability

measure extending  $P$ , it follows that  $P_0$  applied to  $a$  in (5) produces eq.(4), where  $P$  is identified with  $P_0$ .

It is the contention of this paper that a similar product space construction can be used to develop an algebra of numerical and linguistic modifications of probability statements. This is accomplished by generalizing the forms of  $\alpha_f$  in eq.(5), relative to variability of coefficient factors and number of arguments. For example, utilize first the formal relation

$$a^{1/2} = a \bullet (1-a')^{-1/2} \quad (6)$$

and the power series expansion of the second factor, producing the formal power series expansion

$$a^{1/2} = \sum_{j=0}^{+\infty} (a')^j \cdot a \cdot c_j ; \quad c_j = (\text{def}) \binom{2j-1}{j} \bullet 2^{-(2j-1)} , j=0,1,2,\dots \quad (7)$$

In turn, this yields the corresponding product space form

$$a^{1/2} = \bigvee_{j=0}^{+\infty} a' \times \dots \times a' \times a \times \gamma(c_j) , \quad (8)$$

(j factors)

where  $\gamma(c_j)$  is a conditional event such that for all probability measures  $P$  over  $B$ ,

$$P_0(\gamma(c_j)) = c_j . \quad (9)$$

As an example of such *nontrivial constant-valued events*, consider  $\gamma(1/3)$ , representing  $1/3$  in probability for all  $P$  as

$$\gamma(1/3) = (\text{def}) ( b \times b \times b' \mid b \times b \times b' \vee b \times b' \times b \vee b' \times b \times b ) , \quad (10)$$

where  $b \neq \emptyset$ ,  $\Omega$  is any event in  $B$  - to standardize, preferably in limiting form approaching  $\Omega$ , whence

$$\begin{aligned} P_0(\gamma(1/3)) &= P(b)P(b)P(b') / (P(b)P(b)P(b') + P(b)P(b')P(b) + P(b')P(b)P(b)) \\ &= P(b)^2 P(b') / 3P(b)^2 P(b') \\ &= 1/3. \end{aligned} \quad (11)$$

Finally, note that  $a^{1/2}$  is the appropriate appellation for the right-hand side of (8), since  $P_0$  as a product probability measure yields in combination with the disjointness of the terms in eq.(8):

$$P_0(a^{1/2}) = \sum_{j=0}^{+\infty} P(a')^j \cdot P(a) \cdot P(\gamma(c_j)) = P(a) \cdot \sum_{j=0}^{+\infty} P(a')^j \cdot c_j = P(a) \cdot (1 - P(a'))^{-1/2} = P(a)^{1/2}. \quad (12)$$

### References

1. I.R. Goodman, H.T. Nguyen & E.A. Walker, *Conditional Inference and Logic for Intelligent Systems: A Theory of Measure-Free Conditioning*, North-Holland, Amsterdam, 1991.
2. I.R. Goodman, "A new characterization of product space algebra through deductions", to be submitted.
3. I.R. Goodman & H.T. Nguyen, "A theory of conditional information for probabilistic inference in intelligent systems: Part II, product space approach & Part III, mathematical appendix", to appear in *Information Sciences*.